ATTENTION DEFICIT HYPERACTIVITY DISORDER (ADHD)

ReLevant Terms:

- ADD (outdated term): Stood for Attention Deficit Disorder (CDC, 2022)
- ADHD "One of the most common neurodevelopmental disorders of childhood. It is usually first diagnosed in childhood and often lasts into adulthood. Children with ADHD may have trouble paying attention, controlling impulsive behaviors (may act without thinking about what the result will be), or be overly active." (CDC, 2022)
- Neurotypical: "Someone who is neurotypical has a brain that works according to the "normal" standards of our society." (Wake Forest, 2020).
- Neurodivergent: "Neurodiversity is a scientific concept arising from brain imaging. A number of brain studies have shown that people with learning or thinking differences are "wired" differently than their peers. In other words, some children are born with brains that think, learn and process information differently than others." (Wake Forest, 2020).

STUDENT POPULATION: HISTORICAL TRENDS:

- ADHD is a prevalent disability on college campuses. Of students receiving disability services 25% have ADHD (Antshel, 2016)
- These disabilities are hidden and can therefore be affecting anyone from any ethnicity, gender or sexuality
- ADHD is a burdensome, stigmatized, and untreated condition for millions (Barkley, 2014; Doshi et al., 2012; Hinshaw & Scheffler, 2014 as cited in Zhao et al., 2021)
- "Individuals with ADHD are widely stigmatized, as children with ADHD are perceived as more dangerous, lazier, and more shameful than children with asthma, and adults with ADHD are perceived as irresponsible and less socially desirable" (Zhao et al, 2021)
- "Attention Deficit/Hyperactivity Disorder ("ADHD") is a complex multifactorial disorder that was first described in the late 1800s
- In "the early 1900s, ADHD shifted away from being a behavioral based to a neurobiological-based disorder. During this period. individuals with ADHD were classified as having minimal brain damage." (Szeto, 2019)
- Currently treatment options such as psychosocial therapy and medications help ADHD patients maintain a quality of li (Szeto, 2019)
- Evidence-based treatments for ADHD include pharmacological and non-pharmacological therapies (Evans et al. 2014)
 - analysis of the effects of psychosocial treatments over time (including mostly behavioral treatments) shows that in the last 50 years effect sizes for treatments of ADHD have not improved, but rather show a non-significant decline in effectiveness (Weisz et al. 2018 as cited in van der Oord & Tripp,

STUDENTS' COLLEGIATE **EXPERIENCES:**

- Inattentive: Students with ADHD often have trouble in social settings with listening, paying attention and picking up on social cues. (Gill &Hosker, 2021)
- · Hyperactive: Frequently interrupting, Sharing scattered thoughts, Being hyper-focused on a topic, Talking rapidly or excessively. (Gill & Hosker, 2021). Impulsive: Goofy behavior at inappropriate times Entering others' personal space, Displaying aggression, Initiating conversations at inappropriate times. (Gill & Hosker, 2021)
- Combined Type: People with combined-type ADHD demonstrate six or more symptoms of inattention, and six or more symptoms of hyperactivity and impulsivity. (Gill & Hosker, 2021)
- Universal Design: allows students with disabilities to receive accommodations without having to go through their disability services office (Roberson et al., 2022)

Approximate Prevalence Distribution of the Subtypes of ADHD Hyperactive-Impulsive Type Combined Type

Hallmark Signs

of ADHD

SHORT ATTENTION

DISTRACTIBILITY

ORGANIZATIONAL

POOR INTERNAL

PROCRASTINATION

Bowle Terry LEARNING



DIVERSITY WITHIN THE POPULATION:

- Increased undiagnosed cases of ADHD from students with impoverished backgrounds "Low socio-economic status (SES) has been highlighted as a risk factor associated with ADHD among Black individuals. Higher SES is an apparent protective factor for White families but not for Black families. Lower SES has also been tied to familial factors; Black youths are more likely than White youths to be born to young unmarried mothers at birth. Moreover, being a single head of a household, mainly led by single mothers with 4 or more children, is an additional risk factor for Black youths. In addition, Black children who receive a diagnosis of ADHD within a lower SES do not have access to insurance in the US. One study noted that certain families with limited access to insurance do not speak English as a primary language, which serves as a barrier to care. Contrarily, another study found that families living in suburban areas, in contrast to urban areas, have a higher chance of receiving a diagnosis of ADHD due to accessibility to medical care." (Cénat & Blais-Rochette & Morse, 2021).
- Increased undiagnosed cases of ADHD if the student is of color & Less support and accommodation for students of color during youth 'This systematic review and meta-analysis assessing the results of 21 studies found that the pooled prevalence of ADHD among 154818 US Black individuals was approximately 15% and suggested that the associated risk factors included sociodemographic characteristics (age, sex, race, socioeconomic status), risk behaviors, and familial and environmental factors. Meaning The higher prevalence of ADHD observed in this study among US Black individuals highlights the need to increase assessment, accurate diagnoses, and culturally appropriate care.' (Cénat & Blais-Rochette & Morse,

IMPLICATIONS WITHIN HIGHER EDUCATION:

- Accommodations are needed i.e. Testing Time Limits (Extended)
- · Research has shown that if students with ADHD do transfer to higher education, they are more likely to have lower test scores, to repeat classes, and to drop out of higher education without a degree compared with TDC (e.g., Barkley, Fischer, Smallish, & Fletcher, 2006; DuPaul et al., 2009; Kuriyan et al., 2013).
- "Students with diagnosed learning disabilities often receive federally mandated, accommodations in K-12 through a 504 plan, but 504 plans do not exist in the college setting. As such, the support students received in high school may just disappear at college entry".(Kepple, 2021) Typically, students may be invited to share the following:
- a letter disclosing the disability and clarifying current academic impact
- · psychological evaluations diagnosing a disability and the date of diagnosis
- a current IEP/504 plan and records from high school documenting any accommodations and services received (CHADD, 2019)
- Rates of suicide attempts were over four times as high in the ADHD group compared to the non-ADHD group (13.7% vs. 2.9%). (Eddy et al., 2019)

STRATEGIES FOR advising/working with THIS GROUP:

- Identifying specific areas of concern vs. blanketing the issue through 'ADHD Coaching' (Ahmann & Saviet 2021)
- A balanced use of an incentive/consequence system for tasks that need to be completed (Prevatt & Smith 2017)
- Adaptive decompressing methods for feelings of being overwhelmed (Barra & Grub 2021).

